

Study of the nucleon-nucleon bremsstrahlung radiation at COSY-11

P. Moskal, J. Przerwa¹ for the COSY-11 collaboration

The installation of the neutron detector at the COSY-11 facility enables to study a plethora of new reaction channels. It opens wide possibilities not only to investigate the isospin dependence of the meson production [1] but also it enables to measure the bremsstrahlung radiation created in the collisions of nucleons. The study of the latter process is of special interest since it is highly sensitive to the nucleon-nucleon potential, and hence may serve as a tool to discriminate between various existing potential models [2, 3]. The bremsstrahlung radiation may also be helpful for the calibration of the absolute timing of the neutron detector, necessary for the determination of the momentum of neutrons eg. in the analysis of reactions like $pn \rightarrow pn\eta'$. Gamma quants produced in NN collisions are an excellent tool for the calibration of the neutron detector since their velocity is constant and the time-of-flight between the target and the hit module depends on the path length of the γ 's only. Figure 1a presents the time-of-flight spectrum – for the neutral particles – measured between the target and the neutron detector using a detector arrangement as shown in figure 2. The data are from the experiment performed with a deuteron target and a proton beam with a momentum of 2.075 GeV/c [1]. The distribution consists of a rather broad spectrum originating from the neutrons emitted from different reactions, eg. $pn \rightarrow pn\eta$ as shown in figure 1b, and of a sharp peak which we interpret as being due to the γ quants. In the preliminary analysis of the data [1] we have adjusted the absolute time offset of the neutron detector such that the visible peak would correspond to the γ quants. Under this assumption we have obtained satisfactorily coherent results [1], yet naturally, for the final evaluation of the data this assumption must be verified.

At present, in order to perform a more thorough calibration we are analysing data taken in January 2003 from the run with a deuteron beam and proton target. This experiment performed at the beam momentum of 3.204 GeV/c was primarily devoted to study of the $dp \rightarrow dp\eta$ reaction [4], however, we expect that the kinematical conditions when using a deuteron beam and a proton target enabled to register, by the used detection system (fig. 2), many reactions with γ quants in the final state. To study them is interesting in itself, and additionally it will allow us to perform the timing calibration of the whole system.

As a first step we made a presort of the data selecting events with simultaneous signals in any of the drift chambers and the neutron detector. In parallel we have extended the simulation programme to the free and quasi-free nucleon-nucleon reactions with bremsstrahlung radiation. Specifically, we have implemented the free $dp \rightarrow dp\gamma$ and $dp \rightarrow {}^3\text{He}\gamma$ reactions and also the quasi-free reactions $dp \rightarrow d\gamma p_{sp}$, $dp \rightarrow n\gamma p_{sp}$, as well as $dp \rightarrow pp\gamma n_{sp}$. The suffix “ $_{sp}$ ” indicates here the spectator nucleon which does not take part in the reaction. In order to account for the Fermi motion of the nucleons inside a deuteron, we have used an analytical parametrization of its wave function resting from the PARIS potential [5]. Kinematics of events corresponding to the first three of the above listed reactions can be entirely determined by measuring the outgoing charged ejectiles. Therefore, they seem to be best suited for the calibration purposes. However, step by step we will study the response of the COSY-11 de-

tection system to all the above mentioned reactions in order to estimate their usefulness for the calibration of the system and the ability of the determination of their cross sections. Searching for the best calibration method, complementary to the reactions with the bremsstrahlung radiation, we intend also to consider the quasi free proton-proton scattering (see figure 2) and the production of the π^0 meson via the $dp \rightarrow ppn_{sp}\pi^0 \rightarrow ppn_{sp}\gamma\gamma$ process.




Fig. 1: Time-of-flight distribution determined between the target and the neutron detector. The simulation was performed for the $pn \rightarrow pn\eta$ reaction. In addition to neutrons the peak from γ rays is seen at a value of about 25 ns in the experimental spectrum.




Fig. 2: Scheme of the COSY-11 detection system with superimposed tracks from the $dp \rightarrow pp n_{sp}$ reaction. Fast protons are registered in the two drift chambers D1, D2 and in the scintillator hodoscopes. Protons scattered under large angle are measured in the S_{imon} detector. Neutrons are registered in the neutron modular detector (N1), and deuterons in the deuteron detector denoted by DEU.

References:

- [1] P. Moskal et al., e-Print Archive: nucl-ex/0311003.
- [2] N. A. Khokhlov et al., Phys. Rev. C **68** (2003) 054002.
- [3] M. D. Cozma et al., Phys. Rev. C **68** (2003) 044003.
- [4] C. Piskor-Ignatowicz et al., contribution in this report.
- [5] M. Lacombe et al., Phys. Lett. **101 B** (1981) 139.

¹ Institute of Physics, Jagellonian University, Cracow, PL-30-059 Poland